

Evidence Brief:

Inform your practice

Because LGBTQ health matters



Rainbow Health Ontario
Santé arc-en-ciel Ontario

This fact sheet was created
in partnership with:



LGBTQ MENTAL HEALTH

LGBTQ mental health must be understood in the context of historical and ongoing pathologization of LGBTQ identities. With the publication of the DSM-III-R in 1986 the American Psychiatric Association delisted homosexuality as a mental disorder and mental health associations repudiated attempts to change sexual orientation as psychologically damaging.⁽¹⁻²⁾ The World Health Organization removed homosexuality from its list of mental disorders 1990. Despite this, significant stigma is still attached to same-sex attraction in many cultures, and trans people continue to be pathologized as having Gender Identity Disorder, which continues to be listed in the DSM as a diagnostic category.

WHAT IS MENTAL HEALTH?

- The World Health Organization defines mental health as "a state of well-being in which the individual realizes his or her own abilities, can cope with the normal stresses of life, can work productively and fruitfully, and is able to make a contribution to his or her community." LGBTQ mental health is likewise shaped by a sense of self-worth, the level of stress to which people are subjected, and the inclusiveness of their workplaces and community culture.⁽³⁾

MENTAL HEALTH OF LGBTQ PEOPLE

- Large Canadian studies indicate that LGB people are more likely than heterosexuals to report unmet mental health needs and were more likely to consult mental health practitioners.⁽⁴⁾
- Studies have found high rates of depression, anxiety, obsessive–compulsive and phobic disorders, suicidal thoughts and acts, self-harm, and alcohol and drug dependence among LGBTQ people.⁽⁵⁻⁷⁾
- LGB people are one and almost twice as likely to experience childhood maltreatment, interpersonal violence, and personal loss and are at double the risk of developing post traumatic stress disorder as their heterosexual peers.⁽⁸⁾
- Meta-analysis studies found that sexual minority individuals were two and a half times more likely than heterosexuals to have attempted suicide and had a risk of depression and anxiety one and a half times higher than heterosexuals.⁽⁹⁾
- Sexual minority women were particularly at risk for substance-related disorders, while sexual minority men had a higher risk of suicide.⁽¹⁰⁾

- LGBTQ youth have an increased risk of suicide, substance abuse, isolation and experiencing sexual abuse.⁽¹¹⁾ A Canadian study estimated that the risk of suicide among LGB youth is 14 times higher than for their heterosexual peers.⁽¹²⁾ A recent U.S. study of LGBTQ youth found that 10% of them met the criteria for post traumatic stress disorder and 15% met the criteria for major depression.⁽¹³⁾
- A large statistically representative study of trans people in Ontario found that 77% had seriously considered suicide, and 45% had attempted suicide. Trans youth were at greatest risk of suicidality, as were those who had experienced physical or sexual assault.⁽¹⁴⁾

WHY ARE LGBTQ PEOPLE AT HIGHER RISK?

- LGBTQ people experience stigma and discrimination, and this stigma can have a variety of negative consequences throughout the life span.⁽¹⁰⁾ The accumulated stigma, prejudice, and discrimination to which minoritized and marginalized people are exposed is called minority stress.
- LGBTQ people are also the targets of sexual and physical assault, harassment, and hate crimes. The number of hate crimes in Canada motivated by sexual orientation more than doubled from 2007 to 2008, and were the most violent of all hate crimes.⁽¹⁵⁾ These pressures, as well as the stress of sometimes concealing their orientation or modifying their behaviour or appearance in anticipation of homophobia and violence, have a negative effect on mental and physical health.^(9, 16-18)
- A Canadian study suggests that the experience of stigma and discrimination increases internalized homophobia and stress-related cortisol production in LGBTQ people, both of which are associated with increased depression, anxiety and suicidal thoughts.⁽¹²⁾
- LGBTQ people who experience family rejection as adolescents reported high rates of depression, drug use, unprotected sex, and attempted suicide.⁽¹⁹⁾
- Studies suggest that bisexual women may experience higher rates of childhood sexual abuse, childhood physical abuse, intimate partner violence, and non-partner violence than lesbian and heterosexual women.⁽²⁰⁾ Researchers have found connections between these multiple stressors and rates of substance abuse among bisexual women.^(7, 21)
- A study in Ontario found that 20% of trans people had experienced physical or sexual assault due to being trans, and 34% were subjected to verbal threats or harassment.⁽¹⁴⁾
- Trans people in both Canada and the US report high levels of violence, harassment, and discrimination with respect to finding stable housing or employment, and in accessing health or social services.⁽²²⁻²⁶⁾ One US study found that a third of trans people had lost or been denied a job because of their trans identity.⁽²⁷⁾
- Individuals with multiple marginalized identities (such as racialized LGBTQ people) were more likely to report excessive substance use. The prevalence of substance use disorders was more than twice as high among LGB people as it was among heterosexuals.^(7, 21) A study of ethnic minority male-to-female transgender youth found that 18% of them were currently homeless.⁽²⁸⁾
- Poverty may be a factor exacerbating LGBTQ mental health. A Canadian study found that bisexuals were over-represented in the lowest income categories, and an Ontario-based study found that half of trans people were living on less than \$15,000 a year.^(4, 14)

SUPPORTIVE ENVIRONMENTS ARE KEY TO MENTAL HEALTH

- A Canadian Study found that support from family and friends reduced stress and contributed to positive mental health in young gays, lesbians and bisexuals.⁽¹²⁾
- A US study of lesbian and gay parents found that support from family and friends, supportive workplaces and neighbourhoods, and low internalized homophobia promoted positive mental health. Lesbian and gay people with high internalized homophobia, living in states with anti-gay legislation showed the poorest mental health.⁽²⁹⁾
- Experiencing positive responses to coming out is associated with reduced risk of substance abuse.⁽³⁰⁾ Two studies in the US found that 89% of lesbians and bisexual women experienced a negative reaction when they came out to their doctor.^(5, 31) A study of medical students found that one quarter were significantly homophobic, and nine percent viewed homosexuality as a mental disorder.⁽³²⁾
- Youth who identify with the LGBTQ community have been found to significantly reduce their internalized homophobia.⁽³³⁾ Family acceptance of LGBTQ adolescents is associated with good mental and physical health in LGBTQ youth.^(19, 34)

IMPLICATIONS FOR HEALTH CARE PROVIDERS

- Providers should be aware of the broader social and legal context in which their LGBTQ clients live, and explore the degree to which internalized negative social messages may be contributing to their health concerns.
- Suicide response and crisis intervention staff may need additional training to ensure that LGBTQ clients are not subjected to stereotyping or discrimination and that the gender of trans clients is not misidentified.
- Health care providers should be aware of resources available to help those LGBTQ people at greatest risk for suicide. In response to publicized suicides by LGBTQ youth, author Dan Savage initiated the It Gets Better campaign <<http://www.itgetsbetter.org>> in which supportive LGBTQ people and their allies share supportive messages through online video. LGBTQ youth are at high risk for suicide. Youth ages 5-20 can speak with trained counsellors at the Kids Help Phone at 1-800-668-6868. The Lesbian, Gay, Bi & Trans Youthline offers free peer support for youth 26 and under at 1-800-268-9688.
- Given the WHO's definition of mental health, and the effect of minority stress, there should be a recognition that legal and social equity for LGBTQ people is a health issue as well as a political one.
- Promoting family acceptance of LGBTQ adolescents and encouraging them to connect with LGBTQ culture is essential to reducing health disparities among LGBTQ youth. Since family support is protective against stress, PFLAG (Parents, Friends of Lesbians and Gays) <www.pflagcanada.ca> can be a good resource for LGBTQ people and their families.
- Be aware that minority stress may negatively impact physical and mental health minority stress. Be alert to possible social contributors in both mental and physical illnesses. Clients who belong to multiple marginalized communities may face more barriers to maintaining good mental health.

References

- American Psychological Association. Insufficient evidence exists to support the use of psychological interventions to change sexual orientation. 2009.
- American Psychiatric Association. Therapies focused on attempts to change sexual orientation. Arlington, VA: 2000.
- World Health Organization. Promoting mental health: Concepts, emerging evidence, practice: A report of the world health organization. Geneva: Department of Mental Health and Substance Abuse in collaboration with the Victorian Health Promotion Foundation and the University of Melbourne; 2005.
- Tjepkema M. Health care use among gay, lesbian and bisexual Canadians. Statistics Canada. Canada: Statistics Canada; 2008.
- Diamant AL, Wold C. Sexual orientation and variation in physical and mental health status among women. *Journal of Womens' Health*. 2003; 12(1):41-49.
- Cochran SD, Mays VM. Physical health complaints among lesbians, gay men, and bisexual and homosexually experienced heterosexual individuals: Results from the California quality of life survey. *American Journal of Public Health*. 2007; 91(11):2048-2055.
- McCabe S, Bostwick WB, Hughes TL, West BT, Boyd CJ. The relationship between discrimination and substance use disorders among lesbian, gay, and bisexual adults in the United States *American Journal of Public Health*. 2010; 100(10):1946-1952.
- Roberts AL, Austin SB, Corliss HL, Vendermorris AK, Koenen KC. Pervasive trauma exposure among US sexual orientation minority adults and risk of posttraumatic stress disorder. *American Journal of Public Health*. 2010; 100(12):2433-2441.
- King SD. Midlife and older gay men and their use of physical and mental health services: Exploring the effects of health enablers, health need, psychosocial stress and individual health coping. Ohio State University; 2010.
- Meyer IH. Prejudice, social stress, and mental health in lesbian, gay, and bisexual populations: Conceptual issues and research evidence. *Psychological Bulletin*. 2003; 129(5):674-697.
- Gibson P. Gay and lesbian youth suicide. In: Fenleib MR, editor. *The Secretary's Task Force on Youth Suicide, United States Government Printing Report of the Secretary's Task Force on Youth Suicide, United States Government Printing Office*; 1989.
- Benibgui M. Mental health challenges and resilience in lesbian, gay and bisexual young adults: Biological and psychological internalization of minority stress and victimization. 2011.
- Mustanski BS, Garofalo R, Emerson EM. Mental health disorders, psychological distress, and suicidality in a diverse sample of lesbian, gay, bisexual, and transgender youths. *American Journal of Public Health*. 2010; 100(2426-2432).
- Bauer G, Boyce M, Coleman T, Kaay M, Scanlon K, Travers R. Who are trans people in Ontario? Toronto: *Trans PULSE E-Bulletin*; 2010. Report No.: 1(1).
- Dauvergne M. Police reported hate crime in Canada, 2008. *Juristat [Internet]*. 2010; 30(2). Available from: <http://www.statcan.gc.ca/pub/85-002-x/2010002/article/11233-eng.pdf>.
- Pascoe EA, Smart Richman L. Perceived discrimination and health: A meta-analytic review. *Psychological Bulletin*. 2009; 135(4):531-554.
- Selvidge M, Matthews CR, Bridges SK. The relationship of minority stress and flexible coping to psychological well being in lesbian and bisexual women. *Journal of Homosexuality*. 2008; 55(3):450-470.
- American Psychological Association. Answers to your questions: For a better understanding of sexual orientation and homosexuality Washington, D.C.: 2008.
- Ryan C, Russell ST, Huebner D, Sanchez DR. Family acceptance in adolescence and the health of LGBTQ young adults. *Journal of Child and Adolescent Psychiatric Nursing*. 2010; 23(4):205-213.
- Austin SB, Jun HJ, Jackson B, Spiegelman D, Rich-Edwards J, Corliss HL, et al. Disparities in child abuse victimization in lesbian, bisexual, and heterosexual women in the nurses' health study II. *Journal of Women's Health*. 2008; 17(4):597-606.
- Hughes T, McCabe SE, Wilsnack SC, West BT, Boyd CJ. Victimization and substance use disorders in a national sample of heterosexual and sexual minority women and men. *Addiction*. 2010; 105(12):2130-2140.
- Gapka S, Raj R. Trans health project: A position paper and resolution adopted by the Ontario public health association. Toronto: OPHA; 2003.
- Lombardi EL, Van Servellen G. Building culturally sensitive substance use prevention and treatment programs for transgendered populations. *Journal of Substance Abuse Treatment*. 2000; 19:291-296.
- JSI Research & Training Institute, Inc. Access to health care for transgendered persons in greater Boston. Boston: Report for GLBT Health Access Project; 2000.
- Moran LJ, Sharpe AN. Violence, identity and policing. *Criminal Justice*. 2004; 4(4):395-417.
- Namaste V. *Invisible lives: The erasure of transsexual and transgendered people*. Chicago: University of Chicago Press; 2000.
- Xavier J, Simmons R. The Washington transgender needs assessment survey. Washington, D.C.: The Administration for HIV and AIDS of the District of Columbia Government; 2000.
- Garofalo R, Deleon J, Osmer E, Doll M, Harper GW. Overlooked, misunderstood and at-risk: Exploring the lives and HIV risk of ethnic minority male-to-female transgender youth. *Journal of Adolescent Health*. 2006;38:230-236.
- Goldberg E, Smith JZ. Stigma, social context, and mental health: Lesbian and gay couples across the transition to adoptive parenthood. *Journal of Counselling Psychology*. 2011; 58(1):139-150.
- Rosario M, Schrimshaw EW, Hunter J. Disclosure of sexual orientation and subsequent substance use and abuse among lesbian, gay, and bisexual youths: Critical role of disclosure reactions. *Psychology of Addictive Behaviour*. 2009; 23(1):175-184.
- Diamant AL, Schuster MA, Lever J. Receipt of preventive health care services by lesbians. *American Journal of Preventive Medicine*. 2000; 19(3):141-148.
- Klamen DL, Grossman LS, Kopacz DR. Medical student homophobia. *Journal of Homosexuality*. 1999; 37(1):53.
- Cox N, Berghe WV, Dewaele A, Vincke J. Acculturation strategies and mental health in gay, lesbian, and bisexual youth. *Journal of Youth and Adolescence*. 2009; 39(10):1199-1210.
- Ryan C, Hueber D, Diaz RM, Sanchez J. Family rejection as a predictor of negative health outcomes in white and Latino lesbian, gay, and bisexual young adults. *Pediatrics*. 2009; 1:346-352.

ACKNOWLEDGEMENTS

This fact sheet was written by Dr. Margaret Robinson.

*Feedback on this document is welcome. Comments and questions can be addressed to Lorelee Gillis:
lgillis@RainbowHealthOntario.ca*

Last Updated: January 2012